

I. Epidemiological profile

Population	2014	%
High transmission (>1 case per 1000 population)	16 700 000	100
Low transmission (0–1 cases per 1000 population)	0	0
Malaria free (0 cases)	0	0
Total	16 700 000	

Parasites and vectors

Major plasmodium species: *P. falciparum* (100%), *P. vivax* (0%)
 Major anopheles species: *An. funestus*, *An. gambiae*, *An. arabiensis*
 Programme phase: Control

Reported confirmed cases: 2 905 310 Estimated cases, 2013: [2 700 000–4 500 000]
 Reported confirmed cases at community level: 13 523
 Reported deaths: 4490 Estimated deaths, 2013: [2500–11 000]

II. Intervention policies and strategies

Intervention	Policies/strategies	Yes/No	Adopted
ITN	ITNs/LLINs distributed free of charge	Yes	2006
	ITNs/LLINs distributed to all age groups	Yes	2010
IRS	IRS is recommended	Yes	2007
	DDT is authorized for IRS	No	–
Larval control	Use of larval control recommended	No	–
IPT	IPT used to prevent malaria during pregnancy	Yes	1993
Diagnosis	Patients of all ages should receive diagnostic test	Yes	2011
	Malaria diagnosis is free of charge in the public sector	No	–
Treatment	ACT is free for all ages in public sector	Yes	2007
	Sale of oral artemisinin-based monotherapies	Is banned	2011
	Single dose of primaquine is used as gametocidal medicine for <i>P. falciparum</i>	No	–
	Primaquine is used for radical treatment of <i>P. vivax</i>	No	–
	G6PD test is a requirement before treatment with primaquine	No	–
	Directly observed treatment with primaquine is undertaken	No	–
	System for monitoring of adverse reactions to antimalarials exists	Yes	2007
Surveillance	ACD for case investigation (reactive)	No	–
	ACD of febrile cases at community level (pro-active)	No	–
	Mass screening is undertaken	No	–
	Uncomplicated <i>P. falciparum</i> cases routinely admitted	No	–
	Uncomplicated <i>P. vivax</i> cases routinely admitted	No	–

Antimalaria treatment policy

Medicine	Adopted
First-line treatment of unconfirmed malaria	AL 2007
First-line treatment of <i>P. falciparum</i>	AL 2007
Treatment failure of <i>P. falciparum</i>	AS+AQ 2007
Treatment of severe malaria	AS; QN 2007
Treatment of <i>P. vivax</i>	–
Dosage of primaquine for radical treatment of <i>P. vivax</i>	–
Type of RDT used	<i>P. f.</i> only.


Therapeutic efficacy tests (clinical and parasitological failure, %)

Medicine	Year	Min	Median	Max	Follow-up	No. of studies	Species
AL	2005–2012	0	4.45	19.5	28 days	8	<i>P. falciparum</i>
AS+AQ	2005–2012	0	1.7	3.6	28 days	3	<i>P. falciparum</i>

Insecticide susceptibility bioassays (reported resistance to at least one insecticide for any vector at any locality)

Year	Pyrethroid	DDT	Carbamate	Organophosphate	Species/complex tested
2010–2014	Yes	No	Yes	No	<i>An. funestus</i> s.l., <i>An. funestus</i> s.s., <i>An. gambiae</i> s.l.





III. Financing





Financing by intervention in 2014

No data reported for 2014

IV. Coverage



V. Impact



Impact: Insufficiently consistent data to assess trends